

# Coronal Jets

Alphonse C. Sterling

Ronald L. Moore

David A. Falconer

Mitzi Adams

Navdeep Panesar (NPP Postdoc)

Sachiko Akiyama

Seiji Yashiro


Nat Gopalswamy

Francisco Martinez (summer 2016 REU Student)


Supported by NASA's LWS and HGI programs, NASA NPP program, and MSFC/Hinode project.)

# Introduction

- ♦ Coronal jets are well seen in X-rays and in EUV (e.g., Shibata et al. 1992, Shimojo et al. 1994, Cirtain et al. 2007, Nisticò et al. 2009, Raouafi et al. 2016).
- ♦ Often have a “jet bright point” on one side of the jet’s base.
- ♦ Seen in coronal holes, quiet Sun, and active regions.
- ♦ AR jets are similar in appearance to non-AR jets; AR jets are longer and more energetic.



Cirtain et al. (2007)



Sterling et al. (2017)


# Today's Discussion:

- ♦ AR jets are basically the same as non-AR jets; they all fit the "minifilament eruption" magnetic geometry.
- ♦ But, one difference is that frequently a cool minifilament is not apparent in (violent) AR jets.
- ♦ Why not?? (Possible answers provided.)
- ♦ Bonus: What causes minifilament eruptions?


# Coronal Hole Jets

# Coronal Hole Jets: “Minifilament eruptions”

XRT




AIA 193




Sterling, Moore, Falconer, & Adams (Nature,  
2015): 20 Polar CH jets.

(Sterling et al. Event 18

# XRT



# AIA 193




(Sterling et al. Event 3)


# “Normal” Filament Eruption (TRACE)



# XRT




# AIA 211




(Sterling et al. Event 1)

# XRT




# AIA 193




(Sterling et al. Event 2)



# XRT



# AIA 211



(Sterling et al. Event 8)



Sterling et al. (2015, 2016, 2017): “minifilament” eruptions.  
Recently modeled by Wyper, Antiochos, & DeVore (Nature, 2017)

# Quite Sun Jets

# Quiet Sun Jets — Similar to CH jets

AIA 171

AIA 94




(Panesar et al. 2016, ApJL; 10 quiet Sun jets)<sub>13</sub>


# Active Region Jets

(Sterling, Moore, Falconer, Panesar, Akiyama, Yashiro,  
& Gopalswamy ApJ, 2016)


# AIA 171



Sterling et al. (2016, ApJ)




AIA 304



AIA 94

# AIA 171



Sterling et al. (2016, ApJ)

# Active Region Jets: Sterling et al. (2016)

## Results:


- Some AR jets show clear minifilaments; they are slowly developing, less “violent”; surge-like, with weak X-ray signature.
- Other jets show little/no minifilaments; rapidly developing, more violent. Have strong X-ray signature.

# Active Region “Violent” Jets


(Sterling, Moore, Falconer, Panesar, & Martinez  
2017, ApJ)

# AR Jet Example 2: To investigate further, look at a

- different AR — Many violent jets:  
14 Jan 2015 (NOAA AR 12259).
- AIA, HMI, Hinode, IRIS




Hinode/XRT




AIA 193

# HMI on IRIS




Is cool minifilament material hidden by a bright shell?




AIA 193

Bright filament “cocoons” envelope  
some “normal” large-scale eruptions too:




AIA 193

Consider neighboring jetting location:




Hinode/XRT




AIA 193

Minifilament “strand” visible from neighboring region, slightly different time.




AIA 193

What triggers jet-producing minifilament eruptions?




## HMI of IRIS-observed region:




Jets occur at *flux cancelation* locations!

# AR jets (Sterling et al. 2017)




Ave. Cancellation  
rate:  $\sim 10^{19}$  Mx/hr.

Magnetic Evolution with Time




AIA 94 Intensity



Same for QS jets: Occur at cancelation sites.

Ave. Cancelation  
rate:  $\sim 10^{18}$  Mx/hr.



Panesar, Sterling, & Moore (2016) — 10 jets. (Poster!)


# Summary, Discussion, and Conclusions

Detailed investigations of several jets ( $>20$  CH;  $\sim 10$  QS;  $\sim 10$  AR):


- ♦ All occur on neutral lines.
- ♦ Essentially all fit the minifilament-eruption picture (only one or two unclear ones).
- ♦ Some of the less-closely-inspected ones can be unclear, e.g. due to coronal “haze,” small size, or complex magnetic setting.
- ♦ Regarding AR jets:
  - ♦ Slower-buildup ones have obvious erupting minifilaments.
  - ♦ Faster-buildup ones may also have erupting minifilaments, but (1) they may be very thin “strands,” and hence hard to see; (2) cool minifilaments may be hidden by emission (cocoon and/or bright jet spire); (3) etc... (Sterling et al. 2017).
- ♦ AR and QS jets result from episodes of flux cancelation.




AIA 304



AIA 94





# Flux Cancellation Rates:

- For AR jets (~7 events):  $\sim 1.5 \times 10^{19}$  Mx/hr (Sterling et al. 2017).

And from Panesar (2017, private comm.)

- For QS jets (~10 events):  $(1 - 4) \times 10^{18}$  Mx/hr.
- For CH jets (~10 events):  $(2 - 6) \times 10^{18}$  Mx/hr.